



1026 E. 2nd St.
 Casper, WY 82601
 Phone: 307-333-0002
 Fax: 307-333-4425

Esophagogastroduodenoscopy (EGD)

Patient Name: _____ Date of Procedure: _____

___ Casper Surgical Center
 1201 East 3rd St
 PH: (307) 577-2950

___ Summit Medical Center
 6350 E 2nd St
 PH: (307) 232-6600

___ Wyoming Medical Center
 1233 E 2nd St
 PH: (307) 577 -7201

PRIOR TO YOUR PROCEDURE CHECK MEDICATIONS.

If you take any of the following medications, please consult your GI doctor or your primary care provider. Your last dose will be on _____

Aggrenox Pradaxa Xarelto Coumadin Plavix Lovenox Effient Ticlid Jantoven Eliquis

Do not take any of the following medications 7 days prior to your procedure:

Advil	Daypro	Anaprox	Ibuprofen
Arthrotec	Vitamin E	Iron pills	Mobic
Feldene	Relafen	Meloxicam	Piroxicam
Motrin	Aleve	Iodine	Naproxen
Excedrin	Celebrex	Diclofenac	Ketoprofen
Diclofenac	Nabumetone	Naprosyn	Pepto-Bismol

(or any additional NSAIDs not listed above)

**Aspirin 325mg

You may take Tylenol or Acetaminophen for pain.

**If you are taking aspirin under the direction of a cardiologist to prevent a stroke or heart attack, then we will likely have you continue taking it.

Diabetic Instructions

If you are diabetic and on Insulin: Take only ½ of your evening dose and **DO NOT** take the morning dose the day of your procedure.

If you are diabetic and on Oral Medications: **DO NOT** take diabetic medications the evening before or the morning of your procedure.

Morning of procedure

NO SOLID FOODS

- You may have clear liquids from midnight the night before your procedure up until _____ (4 hours prior to your procedure scheduled time) List of clear liquids on back.
- Morning medications may be taken as usual, with the exception of the above listed items.

Check in @ _____ For a _____ procedure time

Please note: Your check in time is subject to change per the request of the surgical center you are scheduled with.

Allowed Clear Liquids	Avoid the following
<p>Clear Liquids Only. We encourage you to drink plenty of the allowed liquids</p> <ul style="list-style-type: none"> • Clear fruit juices without pulp: apple, white cranberry, white grape, lemonade (sweeteners are permitted) • Water, Tea, Black Coffee, clear broth and bouillon (any flavor) • Carbonate soft drinks-regular or diet (Sprite, Ginger Ale, 7-Up, Coke, Pepsi) • Gatorade, Kool-Aid, Jell-O, Ice Popsicles, or other fruit flavored drinks (avoid red, purple, or blue food color) 	<ul style="list-style-type: none"> • Solid Food • Alcohol • Milk or Milk Products • Orange juice

Transportation: You **MUST** make arrangements for someone to drive you home, if you will be receiving any type of anesthesia or sedation. This person must be with you at time of check-in.

If you do not have a ride, your procedure will be cancelled. If you are responsible for young children, make arrangements for their care during your procedure and the remainder of the day.

Revised 7/16/2024